
Air Pollution and Mortality in Chile: Susceptibility among the Elderly

Author(s): Sabit Cakmak, Robert E. Dales, Claudia Blanco Vidal

Source: *Environmental Health Perspectives*, Vol. 115, No. 4 (Apr., 2007), pp. 524-527

Published by: The National Institute of Environmental Health Sciences (NIEHS)

Stable URL: <http://www.jstor.org/stable/4150351>

Accessed: 29/09/2008 16:05

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=niehs>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



The National Institute of Environmental Health Sciences (NIEHS) is collaborating with JSTOR to digitize, preserve and extend access to *Environmental Health Perspectives*.

Air Pollution and Mortality in Chile: Susceptibility among the Elderly

Sabit Cakmak,¹ Robert E. Dales,² and Claudia Blanco Vidal³

¹Department of Statistics, Health Canada, Ottawa, Ontario, Canada; ²Department of Epidemiology, University of Ottawa, Ontario, Canada; ³Area Descontaminacion Atmosferica, Comisión Nacional del Medio Ambiente (CONAMA), Metropolitana De Santiago, Chile

OBJECTIVE: The estimated mortality rate associated with ambient air pollution based on general population studies may not be representative of the effects on certain subgroups. The objective of the present study was to determine the influence of relatively high concentrations of air pollution on mortality in a general population sample and in the very elderly.

STUDY DESIGN: Daily time-series analyses tested the association between daily air pollution and daily mortality in seven Chilean urban centers during 1997–2003. Results were adjusted for day of the week and humidex.

RESULTS: Daily averaged particulate matter with aerodynamic diameter $< 10 \mu\text{m}$ (PM_{10}) was $84.88 \mu\text{g}/\text{m}^3$, sulfur dioxide was 14.08 ppb, and carbon monoxide was 1.29 ppb. The 1-hr maximum ozone was 100.13 ppb. The percentage increases in nonaccidental mortality associated with an increase in PM_{10} equivalent to its mean were 4.53 (t-ratio 1.52) for those < 65 years and 14.03 (3.87) for those > 85 years. Respective values were 4.96 (1.17) and 8.56 (2.02) for O_3 ; 4.77 (2.50) and 7.92 (3.23) for SO_2 ; and 4.10 (2.52) and 8.58 (4.45) for CO.

CONCLUSION: Our results suggest that the very elderly are particularly susceptible to dying from air pollution. Concentrations deemed acceptable for the general population may not adequately protect the very elderly.

KEY WORDS: air pollution, elderly, environment, epidemiology, mortality. *Environ Health Perspect* 115:524–527 (2007). doi:10.1289/ehp.9567 available via <http://dx.doi.org/> [Online 8 January 2007]

Acute exposure to ambient air pollution is associated with increased morbidity and mortality from cardiac and respiratory disease (Stieb et al. 2002). Investigating the possibility of susceptible subgroups is important to better understand the population health effects and guide air quality standards. The very young and very old are easily identifiable subgroups of interest. During the early periods of life, air pollution has been associated with preterm birth, intrauterine growth retardation, sudden infant death syndrome, and infant mortality (Dales et al. 2004; Ha et al. 2003; Liu et al. 2003; Ritz et al. 2002; Wang et al. 1997). Among the elderly, adverse effects of air pollution have been reported in those > 65 years of age (Schwartz 1994), but questions remain. Are these senior citizens more susceptible than those who are younger? What is the effect of air pollution at the further extremes of old age? To address these issues, we examined the association between pollution and mortality during the period 1997–2003 in seven Chilean cities.

Methods

Mortality and pollution data. We classified diseases according to the *International Classification of Disease, 9th Revision* (ICD-9; WHO 1975) codes. We obtained the daily number of deaths in metropolitan Santiago for all nonaccidental (ICD-9 codes < 800), cardiovascular (ICD-9 codes 390–459), and respiratory (ICD-9 codes 460–519) causes from the Instituto Nacional de Estadísticas, the official source of statistical data in Chile. We divided Santiago into urban centers based

on proximity to seven air quality monitoring sites in the following municipalities: Las Condes, Cerrillos, El Bosque, La Florida, Independencia, Santiago Centro, and Pudahuel. The period covered was from 1 January 1997 to 31 December 2003, totaling 2,557 days of observation in each urban center. Each urban center had daily (24-hr mean) measurements of ozone, sulfur dioxide, particulate matter $< 10 \mu\text{m}$ in aerodynamic diameter (PM_{10}), and carbon monoxide throughout the 7-year period of study. Daily average values for PM_{10} , SO_2 , and CO and the daily 1-hr maximum concentrations of O_3 were used in our analysis.

Statistical methods. Daily variations in the number of deaths were related to daily variations in ambient concentrations of air pollutants by a random-effects regression model for count data (Burnett et al. 1995), which assumes a Poisson distribution of daily deaths. Here we assumed a linear association between air pollution and mortality on the logarithmic scale, with the association varying at random between cities. Mortality, air pollution, and weather can vary seasonally and are subject to longer-term time trends. Thus any examination of the association between short-term variations in air pollution and mortality must adjust for temporal trends in mortality and weather factors (Lumley and Sheppard 2003; Ramsay et al. 2003; Samet et al. 2003) and may perform better in this regard than the nonparametric model approaches used in previous studies (Cifuentes et al. 2000).

For each urban center separately, we fit natural spline functions to day of study with

one knot for each of 15, 30, 60, 90, 120, 180, and 365 days of observation. We then selected a model with the number of knots that either minimized the Akaike information criterion (AIC) (Akaike 1973), a measure of model prediction, or maximized the evidence that the model residuals did not display any type of structure, including serial correlation, using Bartlett's test (Priestly 1981). We also plotted model residuals against time and found neither a significant pattern nor correlation. Given our optimal model for time, we then selected our best weather predictor of mortality based on the humidex reading, a measure of the combined effect of temperature and humidity. We considered humidex readings on the day of death and the day before death and accounted for potential nonlinear associations with mortality by using natural spline functions. We examined linear weather models and models with up to 4 knots. The model that minimized the AIC was selected as the optimal weather model for each urban center separately. Indicator functions for day of the week were also included to account for any possible differences in mortality rates among the days of the week.

We then included location-specific time and weather models with air pollution in the regression model. Lagging times of 0–5 days were examined for the air pollutants. We also used unconstrained distributed lags as described by Schwartz (2000). The urban center-specific regression parameter estimates of the pollution–mortality association were then pooled among locations using random-effects models. The pooled estimates of the association between pollution and mortality, estimation of uncertainty, and estimate of variability between urban centers were determined by restricted maximum-likelihood methods (Lindstrom and Bates 1990).

The percentage change in mortality associated with changes in the population-weighted pollutant concentrations, equivalent

Address correspondence to R.E. Dales, The Ottawa Hospital (General Campus), 501 Smyth Rd., Box 211, Ottawa, Ontario, Canada K1H 8L6. Telephone: (613) 737-8198. Fax: (613) 737-8537. E-mail: rdales@ohri.ca

We thank D. Rajs Grzebien of Informacion de Salud, Ministerio de Salud, for providing mortality data, and J. Vera Bascour from Comision Nacional Del Medio Ambiente Region Metropolitana.

The authors declare they have no competing financial interests.

Received 1 August 2006; accepted 8 January 2007.

in magnitude to their mean values, are reported in addition to the ratio of the estimate to its standard error (*t*-ratio). The association between ambient concentrations of air pollutants and mortality was determined for nonaccidental, cardiovascular, and respiratory causes of death, age at death, season, and adjustment for other air pollutants.

Results

Population sizes ranged from 501,000 in the urban center represented by Las Condes to 1.34 million in La Florida, with a total of 5.37 million people in the study based on the 1996 census (Table 1). The death rate varied from 1.03 per 100,000 in Las Condes to 1.97 in Independencia. Total daily deaths averaged 69.69, of which 19.21 were cardiac and 8.38 were respiratory. PM₁₀ concentrations ranged from 65.02 µg/m³ in Las Condes to 91.38 µg/m³ in Pudahuel, with a population-weighted average of 84.88 µg/m³. O₃ concentrations ranged from 84.94 ppb in Pudahuel to 135.52 ppb in Las Condes, with a population-weighted average of 100.13 ppb. SO₂ concentrations ranged from 9.12 ppb in Las Condes to 34.06 in Independencia, with a pooled average of 14.08 ppb; and CO concentrations ranged from 0.92 ppm in Las Condes and to 1.48 ppm in Santiago Centro, with a pooled average of 1.29 ppm.

Pairwise Pearson correlations between pollutants were variable between centers. Correlations between SO₂ and CO and PM₁₀ were always positive, whereas O₃ was negatively correlated with SO₂ and CO in some cities (Table 2). Correlations between PM₁₀ and O₃ and other pollutants varied from negative to positive, -0.55 to 0.82.

The number of knots used for each of the 28 location-pollutant associations was 14, except for associations between O₃ and La Florida, Independencia, and Santiago Centro, for which the number of knots was 42, 84, and 84, respectively. The association between air pollutants and mortality was tested using lags of 0–5 days. In single-pollutant models using the best single-day lag, by urban center, each of the individual pollutants was significantly associated with an increased daily nonaccidental mortality evidenced by a *t*-ratio > 2 (Table 3). There were only two exceptions: PM₁₀ in Las Condes and O₃ in Independencia. Each of the associations were positive and effect sizes varied by < 3-fold between cities.

Averaged over urban centers, the mean single-day lags providing the greatest association were 2 days for O₃, 1 day for SO₂, 1 day for CO, and same day (unlagged) for PM₁₀. With these lags in a single-pollutant model, the percent change in mortality attributable to a change in daily pollutants equivalent in magnitude to their mean ranged from 5.64%

for O₃ to 8.54% for PM₁₀, with *t*-ratios > 2 (Table 4). These effects sizes diminished somewhat but all remained significant after adjustment for the other pollutants.

We examined the mortality effect of air pollution by lag structure, age, cause of death, and season (Table 5). The association between nonaccidental mortality and air pollutants was not significantly different when unconstrained distributed lags were substituted for single-day lags. The distributed but not the single-day effects of air pollution were observed to have an approximate 2-fold greater effect on respiratory than on cardiac deaths (*t*-test, *p* < 0.05 for PM₁₀, SO₂, and CO). Both single-day and unconstrained effects were consistent with an increased susceptibility of dying from air pollution among the elderly. Considering the distributed effects, the percent change in mortality was 2- to 3-fold greater for those at least 85 years of age compared to those < 65 years of age (*t*-test, *p* < 0.05 for PM₁₀, SO₂, and CO).

For single-day lags, age-group differences varied from 3-fold for PM₁₀ to a 57% increase for O₃ (*t*-test, *p* < 0.05 for PM₁₀). Larger risks were observed in the colder months of April–September compared to the warmer period of October–March for PM₁₀ (*t*-test, *p* < 0.05).

Discussion

Kavouras et al. (2001) observed that PM₁₀ concentrations in 1998 in several Chilean cities were high by American and European standards. Our data are similar, with an annual average in metropolitan Santiago of 85 µg/m³ from 1997 through 2003. We did not have concentrations of fine particles < 2.5 µm in diameter (PM_{2.5}) for all centers studied, but expect it would be associated with mortality. In the four where it was available, the annual average ranged from 26 to 35 µg/m³ and the correlations with PM₁₀ varied from 0.78 to 0.90. Ilabaca et al. (1999) also demonstrated a

Table 1. Population, daily mortality, and ambient air pollution concentrations (ppb) by monitoring location.

Region	Population (× 10 ⁵)	Nonaccidental deaths/day	Cardiac deaths/day	Respiratory deaths/day	PM ₁₀ (µg/m ³)	Mean pollutant concentrations		
						O ₃	SO ₂	CO
Las Condes	5.01	5.17	1.61	0.66	65.02	135.52	9.12	0.92
Cerrillos	8.92	11.22	3.01	1.25	84.81	92.94	12.92	1.15
El.Bosque	9.15	11.92	3.15	1.26	86.41	88.02	14.27	1.38
La Florida	13.35	15.52	4.30	1.71	90.22	112.19	13.1	1.43
Independencia	4.21	8.31	2.23	1.10	78.58	89.64	34.06	1.37
Santiago Centro	4.98	8.99	2.83	1.21	82.6	101.72	13.5	1.48
Pudahuel	8.08	8.56	2.08	1.20	91.38	84.94	9.79	1.2
Average	53.70 ^a	69.69 ^a	19.21 ^a	8.38 ^a	84.88 ^b	100.13 ^b	14.08 ^b	1.29 ^b

^aTotal number summed over all seven urban centers. ^bPopulation-weighted average pollutant concentration.

Table 2. Range of Pearson pairwise correlations between pollutants for seven monitoring locations, 1 January 1997 to 31 December 2003.

Pollutant	PM ₁₀	O ₃	SO ₂
O ₃	-0.16 to 0.13		
SO ₂	0.37 to 0.77	-0.04 to 0.78	
CO	0.49 to 0.82	-0.55 to -0.01	0.31 to 0.67

Table 3. Percent change (*t*-ratio) in daily nonaccidental mortality associated with changes in pollutant concentrations equivalent to population-weighted averages by individual location.

Location	PM ₁₀	O ₃	SO ₂	CO
Las Condes	8.37 (1.85)	4.92 (2.31)	9.91 (3.72)	10.58 (3.75)
Cerrillos	11.26 (6.64)	5.80 (3.10)	7.31 (7.30)	5.87 (7.23)
El.Bosque	12.29 (7.33)	6.66 (3.57)	4.84 (5.43)	7.28 (7.96)
La Florida	8.90 (13.08)	3.90 (3.40)	4.15 (4.24)	7.52 (7.92)
Independencia	13.08 (5.42)	2.71 (1.51)	5.98 (5.03)	8.52 (7.89)
Santiago Centro	11.45 (5.83)	5.15 (2.94)	5.48 (5.12)	4.27 (6.06)
Pudahuel	5.77 (3.46)	6.96 (2.66)	6.31 (3.95)	3.48 (4.0)

Single-day lags were used.

Table 4. Pooled city estimates of percent change (*t*-ratio) in daily nonaccidental mortality associated with changes in pollutant concentrations equivalent to population-weighted averages.

Pollutant (mean concentration)	Model adjusted for other pollutants	
	Single-pollutant model	
PM ₁₀ (84.88 µg/m ³)	8.54 (5.14)	6.96 (4.02)
O ₃ (100.13 ppb)	5.64 (2.78)	4.13 (2.00)
SO ₂ (14.08 ppb)	5.65 (4.97)	4.51 (3.27)
CO (1.29 ppm)	5.88 (6.42)	6.13 (4.34)

Single-day lags were used, and values were adjusted for city-specific temporal trends and weather.

strong day-to-day correlation between PM₁₀ and PM_{2.5} between February 1995 and August 1996, with higher median values for both during the colder season. Jorquera et al. (2004) reported that because of geography and climatic conditions, Santiago has a higher ratio of estimated PM₁₀ emissions (tons per year) to annual mean PM₁₀ (micrograms per cubic meter) than Mexico City, Buenos Aires, and São Paulo. A low-lying inversion layer in the winter reduces dispersion of pollutants. Although the present study is based on outdoor-area monitoring, it probably reflects personal exposure. During the winters of 1988 and 1989, Rojas-Bracho et al. (2002) carried out an exposure study of Santiago children 10–12 years of age; personal, indoor, and outdoor PM_{2.5} concentrations were all within 5% at 69.5, 68.5, and 68.1 µg/m³, respectively. Even if the mean outdoor and indoor values are different, our results would be valid as long as the exposures changed in the same direction. The present daily time-series analysis examines the effects of day-to-day differences in air pollution, not absolute values.

Air pollution-related mortality. The present findings averaged over seven urban centers are similar to those of previous air pollution studies in Chile. In 1989 and 1991, cardiac and respiratory mortality were higher on days of increased PM₁₀ (Ostro et al. 1996). Ostro et al. (1996) reported that a 10-µg/m³ change in daily mean PM₁₀ was associated with a 1%

increase in total daily mortality. During the same period, total mortality was associated with PM_{2.5} (Salinas and Vega 1995). Between 1988 and 1996, nonaccidental deaths increased on days of higher air pollution (Cifuentes et al. 2000). Cifuentes et al. (2000) reported that changes in mean levels of pollutants were related to 4–11% changes in mortality. In the present study we found that a change in 24-hr mean PM₁₀ of 10 µg/m³ was associated with a 1% mortality change, using a single-day lag. This effect occurred largely in the colder months when particulate concentrations were higher, with no statistically significant findings in the warmer months. After adjusting for PM₁₀, we also detected an effect of pollutant gases O₃, CO, and SO₂ on mortality. An increase in air pollution was associated with an approximate 50% relative increase in respiratory compared to cardiac deaths for CO, SO₂, and PM₁₀.

Seasonal influences. The mortality effects of CO, SO₂, and PM₁₀ appeared greater during April–September, the colder months, although differences were significant for only PM₁₀. Ilabaca et al. (1999) also reported a seasonal modification of the PM_{2.5} effect on pediatric emergency department visits, greatest in the colder months. In the present study, a change in PM₁₀ of about 85 µg/m³ was associated with a 12.2% change in mortality during the warmer months and 1.3% in the colder months, using unconstrained distributed lags.

During the warmer months, October–March, a change in 1-hr maximum daily O₃ of approximately 100 ppb was associated with a 4.9% change in daily mortality, compared with 2.1% in the colder months. PM_{2.5} is higher in the colder season and O₃ higher in the warmer season. This could affect the estimate of effect if the exposure–response relation was not linear or if there was a threshold effect. However, the pattern of residuals in our analysis was consistent with a linear effect. Threshold effects have not been documented even at levels of air pollution lower than seen in the present study. Another possible reason for seasonal differences may be different time–activity patterns resulting in different exposures. Finally, there is evidence for seasonal differences in particulate mass composition (crustal vs. combustion sourced) and direction (Celis et al. 2004).

Effect modification by age. Bell et al. (2005) reported increased mortality effects in the elderly from O₃ in The National Morbidity, Mortality, and Air Pollution Study of 95 U.S. cities. Bateson and Schwartz (2004) reported that the risk of mortality associated with PM₁₀ in Cook County, Illinois, appeared to increase among elderly women but decreased among elderly men. Filleul et al. (2004) reported a greater effect of air pollution mortality in those > 65 years of age, but it did not reach conventional levels of statistical significance. Others also reported increased susceptibility of those > 65 years of age (Gouveia and Fletcher 2000; Spix et al. 1998). We studied the extremes of old age. Compared with those < 65 years of age, those at least 85 years of age were observed to be over twice as likely to die from acute increases in PM₁₀ and > 50% more likely to die from increases in O₃ and SO₂. Age-related susceptibility was further magnified when unconstrained distributed lags were considered. We also observed a generally monotonic increase in susceptibility with increasingly older age groups. These findings suggest that the determination of air quality guidelines designed to protect the general population may be insufficient to protect the elderly in our society.

In summary, more recent air pollution data in Santiago, Chile, indicate that air pollution levels continues to be high by comparison with those in North America and are associated with stronger mortality effects from respiratory than cardiac disease. Daily increases in gases and particles are associated with increased mortality. The extremely elderly appear to be at greater risk than those who are younger. We recommend that the degree of susceptibility to air pollution in the very elderly be investigated in other countries to determine whether this finding is generalizable across different climates and air pollution characteristics.

Table 5. Percent change (t-ratio) in nonaccidental daily mortality associated with changes in pollutant concentrations equivalent to population-weighted averages by cause of death, age at death, and season.

Classification	PM ₁₀	O ₃	SO ₂	CO
Cause of death				
Nonaccidental				
Single-day lag	8.54 (5.14)	5.64 (2.78)	5.65 (4.97)	5.88 (6.42)
Distributed lag	11.68 (5.22)	4.38 (2.18)	9.28 (6.64)	9.39 (6.89)
Cardiac				
Single-day lag	10.06 (3.25)	8.78 (2.42)	7.24 (3.55)	7.79 (4.56)
Distributed lag	13.33 (3.35)	2.30 (0.78)	10.53 (4.29)	11.22 (4.8)
Respiratory				
Single-day lag	18.58 (4.51)	8.21 (1.46)	12.45 (4.19)	12.93 (5.78)
Distributed lag	29.66 (4.88)	15.63 (2.50)	20.44 (5.21)	21.31 (6.34)
Age at death (years)				
≤ 64				
Single-day lag	4.53 (1.52)	4.96 (1.17)	4.77 (2.50)	4.10 (2.52)
Distributed lag	4.26 (1.29)	1.84 (0.71)	4.27 (2.49)	4.76 (2.19)
65–74				
Single-day lag	9.47 (2.81)	8.00 (1.77)	5.99 (2.49)	6.24 (3.17)
Distributed lag	11.72 (3.01)	2.15 (0.86)	7.21 (2.55)	8.12 (3.88)
75–84				
Single-day lag	12.61 (3.80)	9.42 (2.28)	8.73 (4.00)	8.64 (4.82)
Distributed lag	17.62 (3.72)	3.32 (0.92)	11.2 (4.25)	13.12 (5.12)
≥ 85				
Single-day lag	14.03 (3.87)	8.56 (2.02)	7.92 (3.23)	8.58 (4.45)
Distributed lag	19.73 (3.75)	5.92 (1.92)	11.13 (4.38)	13.20 (4.82)
Season				
April–September				
Single-day lag	9.12 (3.35)	3.21 (1.14)	6.47 (3.92)	7.09 (4.02)
Distributed lag	12.20 (3.75)	2.14 (1.25)	10.23 (4.72)	9.65 (4.50)
October–March				
Single-day lag	0.60 (0.45)	6.19 (1.92)	2.62 (1.19)	5.45 (1.14)
Distributed lag	1.27 (1.46)	4.89 (1.82)	4.25 (1.75)	7.80 (1.89)

REFERENCES

- Akaike H. 1973. Information theory and an extension of the maximum likelihood principle. In: *Second International Symposium on Information Theory* (Petrov BN, Csaki F, eds). Budapest: Akademiai Kiado, 267–281.
- Bateson TF, Schwartz J. 2004. Who is sensitive to the effects of particulate air pollution on mortality? A case-crossover analysis of effect modifiers. *Epidemiology* 15(2):131–132.
- Bell ML, Dominici F, Samet JM. 2005. A meta-analysis of time-series studies of ozone and mortality with comparison to The National Morbidity, Mortality, and Air Pollution Study. *Epidemiology* 16(4):425–426.
- Burnett RT, Ross WH, Krewski D. 1995. Nonlinear random effects regression models. *Environmetrics* 6:85–89.
- Celis JE, Morales JR, Zaror CA, Inzunza JC. 2004. A study of the particulate matter PM₁₀ composition in the atmosphere of Chile. *Chemosphere* 54(5):541–550.
- Cifuentes LA, Vega J, Kopfer K, Lave LB. 2000. Effect of the fine fraction of particulate matter versus the coarse mass and other pollutants on daily mortality in Santiago, Chile. *J Air Waste Manag Assoc* 50(8):1287–1298.
- Dales R, Burnett RT, Smith-Doiron M, Steib DM, Brook JR. 2004. Air pollution and sudden infant death syndrome. *Pediatrics* 113:e628–e631.
- Filleul L, Le Tertre A, Baldi I, Tessier JF. 2004. Difference in the relation between daily mortality and air pollution among elderly and all-ages populations in southwest France. *Environ Res* 94(3):249–253.
- Gouveia N, Fletcher T. 2000. Time series analysis of air pollution and mortality: effects by cause, age and socioeconomic status. *J Epidemiol Community Health* 54(10):750–755.
- Ha EH, Lee J-T, Kim H, Hong Y-C, Lee B-E, Park H-S. 2003. Infant susceptibility of mortality to air pollution in Seoul, South Korea. *Pediatrics* 111:284–290.
- Ilabaca M, Olaeta I, Campos E, Villaire J, Tellez-Rojo M. 1999. Association between levels of fine particulate and emergency visits for pneumonia and other respiratory illnesses among children in Santiago, Chile. *J Air Waste Manag Assoc* 49:154–163.
- Jorquera H, Orrego G, Castro J. 2004. Trends in air quality and population exposure in Santiago, Chile, 1989–2001. *Int J Environ Pollut* 22:507–528.
- Kavouras IG, Koutrakis P, Cereceda-Balic F, Oyola P. 2001. Source apportionment of PM₁₀ and PM_{2.5} in five Chilean cities using factor analysis. *J Air Waste Manag Assoc* 51(3):451–464.
- Lindstrom ML, Bates DM. 1990. Nonlinear mixed effects models for repeated measures data. *Biometrics* 46(3):673–687.
- Liu S, Krewski D, Shi Y, Chen Y, Burnett RT. 2003. Association between gaseous ambient air pollution and adverse pregnancy outcomes in Vancouver, Canada. *Environ Health Perspect* 111:1173–1178.
- Lumley T, Sheppard L. 2003. Time series analyses of air pollution and health: straining at gnats and swallowing camels. *Epidemiology* 14:13–14.
- Ostro B, Sanchez JM, Aranda C, Eskeland GS. 1996. Air pollution and mortality: results from a study of Santiago, Chile. *J Expo Anal Environ Epidemiol* 6(1):97–114.
- Priestly MB. 1981. *Spectral Analysis and Time Series*. London: Academic Press.
- Ramsay T, Burnett R, Krewski D. 2003. The effect of concurrency in generalized additive models linking mortality and ambient air pollution. *Epidemiology* 14:18–23.
- Ritz B, Yu F, Fruin S, Chapa G, Shaw GM, Harris JA. 2002. Ambient air pollution and risk of birth defects in southern California. *Am J Epidemiol* 155:17–25.
- Rojas-Bracho L, Suh HH, Oyola P, Koutrakis P. 2002. Measurements of children's exposures to particles and nitrogen dioxide in Santiago, Chile. *Sci Total Environ* 287:249–264.
- Salinas M, Vega J. 1995. The effect of outdoor air pollution on mortality risk: an ecological study from Santiago, Chile. *World Health Stat Q* 48(2):118–125.
- Samet J, Dominici F, McDermott A, Zeger S. 2003. New problems for an old design: time series analyses of air pollution and health. *Epidemiology* 14(1):11–12.
- Schwartz J. 1994. Air pollution and hospital admissions for the elderly in Detroit, Michigan. *Am J Respir Crit Care Med* 150(3):648–655.
- Schwartz J. 2000. The distributed lag between air pollution and daily deaths. *Epidemiology* 11(3):320–326.
- Spix C, Anderson HR, Schwartz J, Vigotti MA, Letertre A, Vonk JM, et al. 1998. Short-term effects of air pollution on hospital admissions of respiratory diseases in Europe: a quantitative summary of APHEA study results. *Arch Environ Health* 53(1):54–64.
- Stieb DM, Judek S, Burnett RT. 2002. Meta-analysis of time-series studies of air pollution and mortality: effects of gases and particles and the influence of cause of death, age, and season. *J Air Waste Manag Assoc* 52:470–484.
- Wang X, Ding H, Ryan L, Xu X. 1997. Associations between air pollution and low birth weight: A community based study. *Environ Health Perspect* 105:514–520.
- WHO. 1975. *International Classification of Diseases, 9th Revision*. Geneva: World Health Organization.